

# Nicomide®

## Rx Only Dietary Supplement

**DESCRIPTION:** Nicomide® Tablets is a prescription dietary supplement for oral administration, specifically formulated for the dietary management of patients with unique nutritional needs who require increased levels of one or more of the ingredients in this product. Each white-colored, oval-shaped tablet is debossed with “342” on one side and blank on the other.

### Supplement Facts

Serving Size: 1 Tablet

Amount per serving:	% Daily Value	
Niacinamide	750 mg	†
Zinc (as Zinc Bisglycinate Chelate)*	27 mg	180%
Folate (as (6S)-N5-Methyltetrahydrofolic Acid Calcium Salt)	500 mcg	125%
Copper (as Cupric Oxide)	2 mg	100%
Selenium (Selenium Amino Acid Chelate)	50 mcg	71%
Chromium (Chromium Amino Acid Chelate)	100 mcg	83%

† Daily Value (DV) not established.

**OTHER INGREDIENTS:** Methocel K, Lactose Anhydrous, Silicon Dioxide, Stearic Acid, Microcrystalline Cellulose, Magnesium Stearate, Hydroxypropyl Methylcellulose, Triacetin, and Titanium Dioxide.

**INDICATIONS:** Nicomide® Tablets is indicated for use as a dietary supplement for patients who are deficient in, or who are at risk for deficiency in one or more of the ingredients in this product.

**CONTRAINDICATIONS:** This product is contraindicated in patients with a known hypersensitivity to any of the ingredients.

**PRECAUTIONS:** Large doses of Nicomide® Tablets should be administered with caution in patients with a history of jaundice, liver disease, or diabetes. Patients with chronic liver failure and/or renal failure should exercise extreme caution in taking prescribed supplements containing copper.

Folic acid alone is improper treatment of pernicious anemia and other megaloblastic anemias where vitamin B12 is deficient. Folic acid especially in doses above 1.0 mg daily may obscure pernicious anemia in that hematologic remission may occur while neurologic manifestations remain progressive.

**THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.**

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

**WARNINGS:** If you are pregnant, nursing, or taking medication, consult your doctor before use. Use only under the advice and supervision of a physician if you have a history of jaundice, liver disease or diabetes. Abnormal liver function tests have been reported in persons taking daily doses of 500 mg or more of niacinamide. Folate intake should not exceed 250% of the Daily Value (1,000 mcg).

### PREGNANCY & NURSING MOTHERS:

Nicomide® Tablets is not indicated for use as a prenatal/postnatal multivitamin for lactating and nonlactating mothers. Physicians and medical practitioners should administer Nicomide® Tablets with caution to patients who are pregnant, lactating and/or taking medication.

**ADVERSE REACTIONS:** Allergic sensitization has been reported rarely following oral and parental administration of Folate.

**DOSAGE AND ADMINISTRATION:** Take 1-2 tablets daily, as a dietary supplement, as directed by your physician.

**HOW SUPPLIED:** Nicomide® Tablets are white-colored, oval-shaped tablets debossed on one side with “342”, and are supplied in bottles of 60 tablets (75854-343-60). The listed product number is not a National Drug Code. Instead, Avion has assigned a product code formatted according to standard industry practice in order to comply with the formatting requirements of pharmacy and healthcare insurance computer systems.

**STORAGE:** Store at 20° - 25°C (68° - 77°F); excursions permitted to 15° - 30°C (59° - 86°F) [See USP Controlled Room Temperature.]

### MANUFACTURED FOR:

Avion Pharmaceuticals, LLC  
Alpharetta, GA 30022  
1-888-61-AVION Rev. 1117-01

\*Subject to one or more claims of U.S. Patent Nos. 7,582,418, 7,838,042 and 8,425,956.

Nicomide® is a registered trademark of Acella Pharmaceuticals, LLC. Under license from Acella Pharmaceuticals, LLC. All rights reserved.